

The Chow group of a del Pezzo surface over a local field

Chandan Singh DALAWAT

The purpose of this Note is to give complete details of the example sketched in [6] to illustrate the method for calculating the Chow group of a rational surface over a local field split by an unramified extension of the base. The example concerned a del Pezzo surface; the construction of a regular integral model and the determination of the specialisation map are carried out here. As this construction has been referred to in the literature [4], it might not be entirely out of place to make it more widely available.

So let $p \neq 2$ be a prime number, K a finite extension of $\mathbf{Q}_p$, $\mathfrak{o}$ the ring of integers of K and $v : K^\times \rightarrow \mathbf{Z}$ the normalised valuation of K . Let $d \in \mathfrak{o}^\times$ be a unit which is not a square and let $\alpha, \beta, \gamma, \delta \in K^\times$ be invertible elements of the field K whose valuations satisfy $v(\alpha) = v(\beta) + 1$, $v(\delta) = v(\gamma) + 2$. The condition on d (that it be a unit) is essential to the method; the conditions on the valuations of $\alpha, \beta, \gamma, \delta$ are not. They could be related in other ways but the case we have chosen suffices to illustrate the method.

In the projective space $\mathbf{P}_{4,K}$ with coordinates $r : s : t : u : w$, consider the surface defined by :

$$(1) \quad \begin{cases} \alpha(dr^2 - s^2) = \beta(t - u)(t + w), \\ \gamma(dr^2 - t^2) = \delta(s + u)(s + w); \end{cases}$$

it is a del Pezzo surface of degree 4, birational over $K(\sqrt{d})$ to the projective plane. The extension $K(\sqrt{d})$ is *unramified* since d is a unit. So the natural map $\text{Pic } X_{\tilde{K}} \rightarrow \text{Pic } X_{\overline{K}}$ is an isomorphism — where $\overline{K}$ is an algebraic closure of K and $\tilde{K}$ is the maximal unramified extension of K in $\overline{K}$ — and the main technical hypothesis of [6] is satisfied.

In order to apply the method of [6, th. 4], we first need to construct a regular projective $\mathfrak{o}$ -scheme X whose generic fiber X_K is the surface (1) and the irreducible components of whose special fibre are — with their reduced structure — smooth over k , the residue field of K . We will then need to calculate the image of the resulting specialisation map [6, (30)], which amounts to computing the intersection numbers [6, (1)] of the curves in the special fibre X_k with the various irreducible components of X_k , where

$\tilde{k}$ is the residue field of $\tilde{K}$. This is carried out in the following ten steps :

- i). The naïve model X_0
- ii). The first blow-up X_1
- iii). The first special fibre $X_{1,k}$
- iv). Singularities of the transform X_1
- v). The second blow-up X
- vi). The second special fibre X_k
- vii). Regularity of the transform X
- viii). The evaluation map
- ix). The Picard groups
- x). The specialisation map

i).— *The naïve model X_0 .* Let π be a uniformising parameter of K . Multiplying by powers of π and by units, one sees that the surface (1) is the generic fiber of the projective $\mathfrak{o}$ -surface X_0 defined in $E_0 = \mathbf{P}_{4,\mathfrak{o}}$ by :

$$(2) \quad \begin{cases} \pi(dr^2 - s^2) - \beta(t - u)(t + w) = 0, \\ \gamma(dr^2 - t^2) - \pi^2(s + u)(s + w) = 0. \end{cases}$$

Let us determine the singularities of the special fibre $X_{0,k}$ of X_0 , which is given in $\mathbf{P}_{4,k}$ by the system :

$$(3) \quad (t - u)(t + w) = 0, \quad \bar{d}r^2 - t^2 = 0,$$

where $\bar{x} \in k$ denotes the reduction modulo $\pi\mathfrak{o}$ of $x \in \mathfrak{o}$. It has two irreducible components, A_0 and B_0 , defined in $X_{0,k}$ by :

$$A_0 : \quad t + w = 0, \quad B_0 : \quad t - u = 0.$$

The union of the singular loci of A_0 and B_0 is defined in $X_{0,k}$ by the system $r = 0, t = 0$. Since we are interested in constructing an $\mathfrak{o}$ -surface the irreducible components of whose special fibre (with their reduced structure) are smooth over k , let us blow up the closed subscheme of X_0 defined by

$$(4) \quad \pi = 0, \quad r = 0, \quad t = 0.$$

But before doing that, let us see if there are points of $X_{0,k}$, not on (4), which are singular on X_0 : it comes down to checking which points of the intersection $A_0 \cap B_0$, apart from the point $r = 0, t = 0, u = 0, w = 0$ which is already on (4), are in fact singular on X_0 . One verifies that at all these points, with the possible exception of

$$\begin{aligned} R_0 : \quad & \pi = 0, \quad s + t = 0, \quad t - u = 0, \quad t + w = 0, \\ S_0 : \quad & \pi = 0, \quad s - t = 0, \quad t - u = 0, \quad t + w = 0, \end{aligned}$$

A_0 and B_0 are Cartier divisors on X_0 ; as they are regular at these points, so is X_0 . On the other hand, a calculation shows that the two points R_0 and S_0 are singular on X_0 .

ii).— *The first blow-up X_1 .* Let us first construct the ambient space of which X_0 blown up at (4) is going to be a closed subscheme. Let $E_1 = \mathbf{P}(\mathcal{O} \oplus \mathcal{O}(1) \oplus \mathcal{O}(1))$ be the projective bundle over $E_0 = \mathbf{P}_{4,o}$ whose fibres are projective planes $\mathbf{P}_{2,o}$ with coordinates $P : R : T$. The blow-up of X_0 along (4) is by definition the closed subscheme X_1 defined in E_1 by the system (2) and the system

$$(5) \quad \begin{cases} \pi R - rP = 0, \quad \pi T - tP = 0, \quad rT - tR = 0, \\ \gamma(dR^2 - T^2) - P^2(s + u)(s + w) = 0. \end{cases}$$

iii).— *The first special fibre $X_{1,k}$.* The special fibre $X_{1,k}$ of X_1 is defined in $E_{1,k}$ by the system (3) and the reduction modulo π of the system (5). The four irreducible components of $X_{1,k}$ are defined in X_1 by the systems

$$\begin{aligned} A_1 : \quad & P = 0, \quad t + w = 0, & C_1 : \quad & \pi = 0, \quad t = 0, \quad u = 0, \\ B_1 : \quad & P = 0, \quad t - u = 0, & D_1 : \quad & \pi = 0, \quad t = 0, \quad w = 0. \end{aligned}$$

It is to be noted that, k' being the quadratic extension of k obtained by adjoining a square root of $\bar{d}$, there is natural morphism $A_{1,k'} \rightarrow \mathbf{P}_{2,k'}$ (coordinates $r : s : u$) whose fibre at a given closed point x is the pair of points $P = 0, \bar{d}R^2 - T^2 = 0$ of the projective plane $\mathbf{P}_{2,k'}$ (coordinates $P : R : T$) which is the fibre of $E_1 \rightarrow E_0$ at x . The irreducible component A_1 thus consists of two conjugate projective planes which do not meet; in particular, it is smooth over k . The same holds for B_1 .

As for C_1 , one sees that the fibre of the projection $C_1 \rightarrow \mathbf{P}_{1,k}$ (coordinates $s : w$) at a given point $(s : w)$ is the conic

$$\bar{\gamma}(\bar{d}R^2 - T^2) - P^2s(s + w) = 0$$

in the plane $\mathbf{P}_{2,k}$ fibre of $E_1 \rightarrow E_0$ at the said point. Thus, C_1 is a conic bundle over $\mathbf{P}_1$ with two degenerate fibres, namely those above $(s : w) = (0 : 1)$ and $(1 : -1)$; in particular, C_1 is smooth and absolutely irreducible. The same holds for D_1 , which is a conic bundle over $\mathbf{P}_{1,k}$ (coordinates $s : u$) whose fibre at any given point $(s : u)$ is

$$\bar{\gamma}(\bar{d}R^2 - T^2) - P^2(s + u)s = 0.$$

It should also be noted that the intersection of A_1 and B_1 is a pair of conjugate lines which do not meet and that C_1 and D_1 meet in the conic

$$(6) \quad \bar{\gamma}(\bar{d}R^2 - T^2) - P^2s^2 = 0$$

in the fibral $\mathbf{P}_{2,k}$ at the point of intersection of the two lines above which C_1 and D_1 are conic bundles.

iv).— *Singularities of the transform X_1 .* Let us determine the singularities of the strict transform X_1 (given by the systems (2), (5)). As the projection $X_1 \rightarrow X_0$ is an isomorphism in the complement of $C_1 \cup D_1$ in X_1 , the inverse images of R_0 and S_0 , namely

$$\begin{aligned} R_1 : \quad & P = 0, \quad s + t = 0, \quad t - u = 0, \quad t + w = 0, \\ S_1 : \quad & P = 0, \quad s - t = 0, \quad t - u = 0, \quad t + w = 0, \end{aligned}$$

are singular on X_1 . All other singularities of X_1 , if any, must lie on $C_1 \cup D_1$.

Now, at every point of $C_1 \cup D_1$ not in their intersection $C_1 \cap D_1$, the divisors A_1, B_1, C_1, D_1 are Cartier; since they are also regular, so is X_1 at those points. The trace of $C_1 \cap D_1$ on the open subscheme $Y : P \neq 0, s \neq 0$ of X_1 is defined by the ideal $\mathcal{I}$ generated by π, u and w . The $(\mathcal{O}_Y/\mathcal{I})$ -module $\mathcal{I}/\mathcal{I}^2$ is free of rank 2 : one has the relation

$$\pi = -\beta(t - u)(t + w) + \pi dr^2$$

which shows that $\pi \in \mathcal{I}^2$. As the trace in question (the conic (6) deprived of its closed point $P = 0$) is regular of codimension 2, it follows that X_1 is regular at these points.

Finally, it turns out that the closed point defined in $C_1 \cap D_1$ by $P = 0$ and in X_1 by

$$M_1 : \quad P = 0, \quad t = 0, \quad u = 0, \quad w = 0$$

is indeed singular on X_1 ; it is also the point where the four irreducible components of $X_{1,k}$ meet. Thus the only singularities of X_1 are the three points R_1, S_1, M_1 ; their union is defined by

$$(7) \quad P = 0, \quad t - u = 0, \quad t + w = 0, \quad t(dr^2 - s^2) = 0.$$

v).— *The second blow-up X .* We first construct an ambient space E_2 from E_1 in the same manner as E_1 was constructed out of E_0 in *ii)* above, but this time the fibres are $\mathbf{P}_{3,0}$ (coordinates $V : U : W : Z$) of “weights” $(0 : 1 : 1 : 3)$ (just as they were of weights $(0 : 1 : 1)$ in *ii)* above). In E_2 , the blow-up X of X_1 along (7) is defined by the systems (2), (5) and the system

$$\begin{cases} \text{rk} \begin{pmatrix} V & U & W & Z \\ P & t - u & t + w & t(dr^2 - s^2) \end{pmatrix} < 2, \\ VZ - \beta T.UW = 0. \end{cases}$$

vi).— *The second special fibre X_k .* The seven irreducible components of the special fibre X_k are the following :

- A : the blow-up of A_1 at the closed points R_1, S_1, M_1 ,
- B : the blow-up of B_1 at the closed points R_1, S_1, M_1 ,
- C : the blow-up of C_1 at the closed point M_1 ,
- D : the blow-up of D_1 at the closed point M_1 ,
- R : the fibre above the closed point R_1 ,
- S : the fibre above the closed point S_1 ,
- M : the fibre above the closed point M_1 .

where fibres of the projection $X \rightarrow X_1$ are meant. All seven irreducible components (with their reduced structure) are smooth over k . The first six are of multiplicity 1; let us show that M is of multiplicity 2. The functions T and $dr^2 - s^2$ are invertible on M and $t - u$ is a local equation for M in the open subscheme of X where T, $dr^2 - s^2$ and U are invertible. In this open subscheme, we have $\pi = \frac{tP}{T}$, $t = \frac{Z}{(dr^2 - s^2)U} \cdot (t - u)$ and $P = \frac{V}{U} \cdot (t - u)$, from which we get the relation

$$\pi = \frac{VZ}{T(dr^2 - s^2)U^2} \cdot (t - u)^2,$$

showing that M is of multiplicity 2 in X_k . The divisor corresponding to the special fibre thus consists of

$$(8) \quad \begin{array}{ccccccc} A & B & C & D & R & S & 2M \\ 2 & 2 & 1 & 1 & 2 & 2 & 2 \end{array}$$

where the second row indicates that the only absolutely irreducible components are C and D, each of the five others splits as two irreducible components over $\tilde{k}$ — the residue field of $\tilde{K}$. These are in fact defined over the quadratic extension of k in which $\bar{d} \in k$ is a square.

The mutual intersections of these seven components are easy to compute and will be of use later.

vii).— *Regularity of the transform X.* Let D be the union of the three divisors R, S, M of X; its complement is regular since it is isomorphic to X_1 deprived of the closed subscheme (7), which we know to be regular (cf. iv) above). Now the Cartier divisor D (defined by $P = 0$ when $V \neq 0$, by $t - u$ when $U \neq 0$, by $t - w$ when $W \neq 0$ and by $t(dr^2 - t^2)$ when $Z \neq 0$) is regular and hence so is X at each of its points. This shows that the scheme X is everywhere regular.

viii).— *The evaluation map* [6, p. 7]. Let F be the free $\mathbf{Z}$ -module on the irreducible components of $X_{\tilde{k}}$ and let $(f_Y)_Y$ be the canonical basis — indexed by the irreducible components of X_k — of $\text{Hom}_k(F, \mathbf{Z})$ (where Hom_k stands for $\text{Gal}(\tilde{k}|k)$ -homomorphisms). We have the evaluation map $\xi : \text{Hom}_k(F, \mathbf{Z}) \rightarrow \mathbf{Z}$ which to a $\text{Gal}(\tilde{k}|k)$ -homomorphism $F \rightarrow \mathbf{Z}$ associates its value at the special fibre $X_{\tilde{k}}$ considered as an element of F (cf. (8)). From what we have said about the multiplicities of the components and about their absolute irreducibility, one has

$$(9) \quad \begin{array}{cccccccc} Y = & A & B & C & D & R & S & M \\ \xi(f_Y) = & 2 & 2 & 1 & 1 & 2 & 2 & 4. \end{array}$$

ix).— *The Picard groups*. In order to complete the calculation of the Chow group of the K-surface (2) — the generic fibre of X — by the method of [6, th. 4], we wish to determine the Picard groups of the seven irreducible components of X_k .

— Pic A. Let $\mathfrak{l}_A$ be the inverse image under $A \rightarrow A_1$ of a pair of conjugate lines in A_1 not passing through the points R_1, S_1, M_1 . Let $\mathfrak{d}_{R,A}$ be the inverse image of a pair of conjugate lines in A_1 passing through R_1 but distinct from the pair $A_1 \cap B_1$. Similarly, define $\mathfrak{d}_{S,A}$ (resp. $\mathfrak{d}_{M,A}$) to be the inverse image of a pair of conjugate lines in A_1 passing through S but distinct from $A_1 \cap B_1$ (resp. passing through M but distinct from $A_1 \cap D_1$). The group $\text{Pic } A$ is generated by the classes of the four divisors $\mathfrak{l}_A, \mathfrak{d}_{R,A}, \mathfrak{d}_{S,A}, \mathfrak{d}_{M,A}$.

— Pic B. In the preceding discussion, if we replace A by B , B by A , C by D and D by C , we get four divisors $\mathfrak{l}_B, \mathfrak{d}_{R,B}, \mathfrak{d}_{S,B}, \mathfrak{d}_{M,B}$ whose classes generate the group $\text{Pic } B$.

— Pic C. Let $\mathfrak{f}_C$ be the inverse image under $C \rightarrow C_1$ of a fiber of $C_1 \rightarrow \mathbf{P}_1$ which does not pass through the point M_1 . The group $\text{Pic } C$ is generated by the classes of the three divisors $\mathfrak{f}_C, C \cap B, C \cap M$.

— Pic D. In an analogous fashion, the group $\text{Pic } D$ is generated by the classes of the divisors $\mathfrak{f}_D, D \cap A, D \cap M$.

— Pic R, Pic S, Pic M. The group $\text{Pic } R$ is generated by the divisors $R \cap A$ and $R \cap B$. The same is true for $\text{Pic } S$ and for $\text{Pic } M$.

x).— *The specialisation map* [6, p. 7]. We need to determine the images of the specialisation maps $\text{Pic } Y \rightarrow \text{Hom}_k(F, \mathbf{Z})$ [6, (30)], where Y runs through the irreducible components of X_k . In other words, we have to determine the subgroup of $\text{Hom}_k(F, \mathbf{Z})$ generated by the images of the various divisors on X_k which we have listed in ix) above.

But consider for example the *divisor* $C \cap B$ on C . Its *image in* $\text{Pic } B$ is a linear combination of the classes of $\mathfrak{l}_B, \mathfrak{d}_{R,B}, \mathfrak{d}_{S,B}, \mathfrak{d}_{M,B}$ and hence its image in $\text{Hom}_k(F, \mathbf{Z})$ is contained in the image of $\text{Pic } B$.

Ruling out such cases, we are reduced to computing the images of the ten divisors which have had the honour of being designated by a gothic letter. These images are expressed in terms of the canonical basis $(f_Y)_Y$ of $\text{Hom}_k(F, \mathbf{Z})$, where Y runs through the irreducible components (8) of the special fibre X_k :

$$\begin{array}{c} f_A \\ f_B \\ f_C \\ f_D \\ f_R \\ f_S \\ f_M \end{array} \begin{pmatrix} \mathfrak{l}_A & \mathfrak{d}_{R,A} & \mathfrak{d}_{S,A} & \mathfrak{d}_{M,A} & \mathfrak{f}_D & \mathfrak{l}_B & \mathfrak{d}_{R,B} & \mathfrak{d}_{S,B} & \mathfrak{d}_{M,B} & \mathfrak{f}_C \\ -2 & -1 & -1 & -2 & 1 & 1 & -2 & -1 & -1 & -2 & 1 \\ 1 & & & & & -2 & -1 & -1 & -2 & & -2 \\ 2 & & & & -2 & & & & & & \\ & 1 & & & & & 1 & & & & \\ & & 1 & & & & & 1 & & & \\ & & & 1 & & & & & 1 & & \end{pmatrix}$$

Notice that the “degree” of each column is 0 (*i.e.* ξ vanishes at that element of $\text{Hom}_k(F, \mathbf{Z})$), in accordance with [6, lemma 10] — for example the degree of the first column is

$$\xi(-2f_A + f_B + 2f_D) = (-2).2 + 1.2 + 2.1 = 0$$

(cf. (9)). Now the map γ_0 [6, (16)] is an isomorphism of the Chow group $A_0(X_K)_0$ of 0-cycles of degree 0 on X_K with the torsion subgroup of the quotient of $\text{Hom}_k(F, \mathbf{Z})$ by the subgroup generated by the columns of the displayed matrix [6]. Hence :

PROPOSITION. — *The Chow group $A_0(X_K)_0$ of 0-cycles of degree 0 on the surface (2) is isomorphic to $\mathbf{Z}/2\mathbf{Z}$.*

The result is in conformity with Coombes and Muder [2, th. 4.5], who used methods specific to del Pezzo surfaces (split by an unramified extension of the base).

If we use Coray and Tsfasman [3, prop. 4.6] to write (2) as the Châtelet surface

$$(10) \quad y^2 - dz^2 = x(x - e_1)(x - e_2), \quad e_1 = \gamma - \pi^2 + \frac{\pi^3}{\beta}, \quad e_2 = \frac{\gamma\pi}{\beta},$$

the result agrees with a calculation of Colliot-Thélène [3, prop. 4.7] (see also [5, prop. 1]) as the Chow group is a birational invariant [1, prop. 6.3].

Let us terminate with a word about the case when $d \in K^\times$ is not a unit in (1), (10); the extension $K(\sqrt{d})$ of K is then *ramified*. No general method seems to be available for calculating the Chow group of a rational surface over a local field which is not split by an unramified extension of the base. But in this particular case, the Chow group has been calculated in [5, prop. 2] as a function of the type of bad reduction (cf. [5, prop. 5]).

BIBLIOGRAPHICAL REFERENCES

- [1] COLLIOT-THÉLÈNE (J.-L.) et CORAY (D. F.). — *L'équivalence rationnelle sur les points fermés des surfaces rationnelles fibrées en coniques*, Compositio Math., **39** (3), 1979, p. 301–332
- [2] COOMBES (K. R.) and MUDER (D. J.). — *Zero-cycles on del Pezzo surfaces over local fields*, Journal of Algebra **97** (1985), 438–460.
- [3] CORAY (D.) and TSFASMAN (M. A.). — *Arithmetic on singular del Pezzo surfaces*, Proceedings of the London Math. Soc. (3) **75** (1988), 25–87.
- [4] CORTI (A.). — *Del Pezzo surfaces over Dedekind schemes*, Ann. of Math. (2) **144** (1996) 3, 641–683.
- [5] DALAWAT (C. S.). — *Le groupe de Chow d'une surface de Châtelet sur un corps local*, Indag. mathem. N.S. **11** (2) (2000), 173–185, [math.AG/0302156](#).
- [6] DALAWAT (C. S.). — *Le groupe de Chow d'une surface rationnelle sur un corps local*, [math.AG/0302157](#).

Chandan Singh Dalawat
 Harish-Chandra Research Institute
 Chhatnag Road, Jhansi
 ALLAHABAD 211 019 India